



Isolation, characterization and identification of microorganisms capable of growing in chocolate pralines

Nielsen, Cecilie Lykke Marvig; Thorsen, Line; Nielsen, Dennis Sandris

Publication date:
2012

Document version
Peer reviewed version

Citation for published version (APA):
Nielsen, C. L. M., Thorsen, L., & Nielsen, D. S. (2012). *Isolation, characterization and identification of microorganisms capable of growing in chocolate pralines*. Poster session presented at International ICFMH Symposium FoodMicro 2012, Istanbul, Turkey.

C.L. Marvig, L. Thorsen, D.S. Nielsen
University of Copenhagen, Faculty of Science, Department of Food Science, Denmark
clmn@life.ku.dk

Chocolate pralines can be spoiled by growth of microorganisms in the filling changing the sensory appearance of the product by production of off-flavors and through crack formation due to gas production. Chocolate praline filling contains 30-50% sugar and up to 10% alcohol, which eliminates growth of most microorganisms. Nonetheless, microorganisms tolerating low water activity have previously been shown to grow in high sugar content food products.



Characterization: MRS broth with 40-70% sucrose and 0-12% ethanol. Growth measured as OD at 460nm.

Samples with high CFU/g counts contained mainly yeast. However, bacteria was found in counts up to 10^3 CFU/g.



Off-flavors, gas, and slime was produced.



- On media modified with 40% sucrose, 10^7 CFU/g and 10^5 CFU/g was isolated from chocolate pralines and production environment, respectively.
- *Staphylococcus* spp., *Bacillus* spp., *Zygosaccharomyces* spp., *Candida* spp., *Wickerhamomyces* spp., *Citeromyces* spp., *Aspergillus* spp., and *Penicillium* spp. was isolated.
- *Wickerhamomyces anomalus* and *Zygosaccharomyces rouxii* grow under conditions relevant for spoilage of chocolate pralines.